

# Prevalence of Diabetes Mellitus in a Tertiary Health Institution in Gombe Metropolis, Nigeria

Mela Danjin<sup>1</sup>, Nafisa Usman Umar<sup>2</sup>, Danladi Adamu<sup>3</sup>

ORIGINAL ARTICLE

## ABSTRACT

**Aim:** This study sought to determine the prevalence of diabetes mellitus (DM) among adult patients attending Specialist Hospital Gombe, Northern Nigeria.

**Materials and Methods:** A 5-year (2010-2014) retrospective descriptive survey of cases of DM was conducted, using hospital record retrieved from the record (Institution). Diagnosed adult cases of DM were carefully identified and sorted from the general outpatient department morbidity register. Relevant data were then harvested on a specially designed pro forma/data collection template.

**Results:** A total of 40,083 medical cases were counted out of which 2069 cases of DM were identified. This implies an overall prevalence rate of 5.16%. However, the yearly prevalence of the disease revealed as high as 12.55% in the year 2012 and a least value of 1.22% in the year 2010. Age-specific case count of the disease was found to be highest, 1288 (62.25%) within the highest age range of 65-75 while the lowest age range of 25-34 was least affected, 33 (1.60%). Overall, more females, 1491 (72.06%) suffered from the disease than males, 578 (27.94%). The sex difference was found to be significant ( $\chi^2 = 449.15521$ ,  $P < 0.001$ ). Furthermore, the case burden of the disease was found to be significantly ( $\chi^2 = 39.02524$ ,  $P < 0.001$ ) highest among individuals living in urban areas, 663 (80.38%) than those living in rural areas, 406 (19.62%).

**Conclusion:** The disproportionately high case burdens of DM among females, urban dwellers and the aged are useful information for the design of targeted diabetes prevention programs by government and other stakeholders.

**KEY WORDS:** Adult patients, diabetes, prevalence, retrospective.

## Introduction

In developing countries such as Nigeria, diabetes mellitus (DM), and other non-communicable diseases (NCDs) continue to exhibit the “slow epidemic” surge. Based on the 2013 IDF global diabetes scorecard, raw national diabetes prevalence for Nigeria was estimated at 4.99%.<sup>[1]</sup> The national figure for some other African countries was put at 3.58% for Kenya, 3.93% for Guinea, 3.35% for Ghana, and 1.55% for The Gambia. Ethiopia had 4.36%, DR Congo 5.37%, Rep of Congo 5.48%, and Cameroun 4.88%.<sup>[1]</sup> From Abeokuta, South Western Nigeria, an overall prevalence rate of 14.8% was reported.<sup>[2]</sup> When segregated by age group an increasing trend

from 2.9% in the age group of 18-27 years to a peak of 21.1% in the age group of 48-57 years was observed. Omorogiuwa *et al.* in a survey carried out on a university community, Ekpoma, Edo State southern Nigeria; reported an overall prevalence of as high as 9.8%.<sup>[3]</sup> In Umudike, Abia State South Eastern Nigeria in a study among adult Nigerians 18 years and above, a prevalence rate of 3.0% with higher rates reported among females (3.6%) than males (2.3%) was published.<sup>[4]</sup> In another study, among adults aged 18 years and above residing in five rural communities in Edo state an overall crude prevalence of 4.6% with sex differences (males 1.9%, females 5.8%,  $P = 0.012$ ) was reported.<sup>[5]</sup> The overall prevalence of as high as 10.5% with male and female rates of 9.6% and 11.2%, respectively, was reported in Uyo, South Southern Nigeria.<sup>[6]</sup>

In northern Nigeria, the prevalence and incidence surveys also indicate varying figures. A 5-year (2001-2005) retrospective study of hospital records in Katsina was conducted in which among the

### Access this article online

Quick Response Code:



Website: [www.jmsh.ac.in](http://www.jmsh.ac.in)

<sup>1</sup>Head of Department, Department of Anatomy and Physiology, College of Nursing and Midwifery, Gombe, Nigeria,

<sup>2</sup>Student, College of Nursing and Midwifery, Gombe, Nigeria, <sup>3</sup>Principal, School of Nursing and Midwifery Gombe, Nigeria

#### Address for correspondence:

Mela Danjin, Department of Anatomy and Physiology, College of Nursing and Midwifery, Gombe, Nigeria.

Phone: +2348063583988. E-mail: danjin67@yahoo.com

281 cases reported in the period, more males (201 [71.53%]) suffered from the disease than females (80 [28.47%]).<sup>[7]</sup>

Although this study did not aim to segregate DM cases by type, extent data from other African countries and Nigeria reveals that about 90% of cases are Type 2 DM while Type 1 DM ranges between 5.7% and 10.0% with the remaining 0.0-4.3% being gestational DM.<sup>[8-14]</sup>

Despite the fact that there is no published work, anecdotal evidence has indicated high burden of DM in the study area. A cursory look at the morbidity register of State Specialist Hospital Gombe would show that DM is a disease of public health importance in the area. Hence, this study was an attempt to document scientific evidence and quantification of the problem within Gombe metropolis. The study also sought to find out trend of DM case prevalence in retrospect (2010-2014) and its seasonal pattern of occurrence. Again, what were the age group, gender, and place of residence patterns of the disease distribution among the study population?

## Materials and Methods

### Study setting

This survey was conducted in the State Specialist Hospital, domiciled within Gombe metropolis which is the state capital of Gombe state in northern Nigeria. The Metropolis lies between latitude 11°08'1 N and 11°02'41 N, longitude 11°02'21 E and 11°01'18 E. The temperature averages 30°C with an annual rainfall of 52 cm. The hospital is the only state government-owned tertiary health facility that is meant to cater for referral cases from other secondary (General Hospitals) and primary health facilities scattered over the 11 local governments of the state. Hence, to a large extent, the hospital patient demographics may be a fair reflection of the entire state. Some of the specialist units or departments include general outpatient department which serves as a filter where all cases are first seen, Paediatrics Department, a Standard Laboratory, Eye Clinic, Dental Clinic, Obstetric and Gynaecology Department, Health Records Department, etc. Based on the Nigerian National Population Commission Census 2006 figures projected to 2014, Gombe state has an estimated total population of about three Million at an annual growth rate of 3.2%.<sup>[8]</sup> The metropolis has a projected population of 344,804 inhabitants.

### Study design

A retrospective descriptive survey of DM was conducted in specialist hospital where the records in the health record department of the hospital were retrieved, reviewed, and analyzed with the aim of determining overall prevalence of the disease as well as characterizing trends and patterns of its distribution based on some selected demographics of the patients.

### Target population

#### Inclusion criteria

To be included in the survey, a case record must be that of an adult patient seen in the outpatient department of Specialist Hospital Gombe with medical diagnosis of DM within the period of 2010-2014; and must be within the age range of 25-74 years (well over 90% of all DM cases retrieved). The record of all other medical cases seen within the same period qualified for the overall all-cases count.

#### Exclusion criteria

A case record will be excluded especially if patient was seen before the year 2010 or after the year 2014. Only very few diagnosed DM cases fell outside the age range of 25 and 75 years. For the general all-cases count cases outside the study period (2010-2014) were excluded from the study.

### Data collection

The relevant hospital records of all medical cases including DM in the state (institute), from 2010 to 2014, were carefully sorted with the aim of investigating the prevalence of the DM in Gombe. A total of 40,083 medical cases were counted, out of which 2069 cases of DM were identified. The key variables on which data were collected include; number of all medical cases seen, number of cases of DM seen, age range, sex, place of residence, month and year.

A structured pro forma with columns under the aforementioned key variable headings was designed and used for the data collection on all identified diagnosed DM cases.

### Data analysis

The raw data were entered on Microsoft excel spreadsheet for preliminary analysis. Thereafter, it was exported to Epi Info 7 (CDC, Atlanta, GA, USA, 2011) for further analysis. Data were presented as frequency tables, bar charts, and graphs. Associations were tested using chi-square within 95% confidence interval (CI).


**Ethical consideration**

Ethical clearance for this work was obtained from the Research and Ethics Committee of College of Nursing and Midwifery Gombe, while consent of relevant authorities of the hospital was properly secured.

**Results**

The results of this study showed a total number of all medical cases seen in the hospital within the study period (2010-2014) as 40,083 out of which 2069 were cases of DM (Table 1). This represented an overall DM prevalence of 5.16%. The yearly trend of DM prevalence is also shown in Table 1, 2010 (1.22%; 76/6212), 2011 (3.96%; 226/5704), 2012 (12.55%; 1219/9710), 2013 (3.33%; 431/12944), and 2014 (2.12%; 117/5513). Table 2 shows DM cases seen within the study period according to age group as, 25-34 (33; 1.60%), 35-44 (127; 6.14%), 45-54 (162; 7.83%), 55-64 (459; 22.18%), and 65-75 (1288; 62.25%). The increases in the disease burden with age was significant ( $\chi^2 = 24.5502, P = 0.000002$ ). In Table 3, sex distribution of DM cases within the study period was shown with significant

( $\chi^2 = 449.15521, P = 0.000000$ ) female majority, male (578; 27.94%) and female (1491; 72.06%). Figure 1 displays the cumulative monthly trend of the 2069 DM cases identified within the 5-year study period (2010-2014). The lowest case count (54) was in the month of March, while the three peaks were recorded in the months of July (141), October (428), and December (560). Table 4 shows DM case burden according to place of residence in which 80.38% (1663/2069) of the patients domiciled in urban areas



**Figure 1:** Monthly trend of diabetes mellitus case count in specialist hospital Gombe (2010-2014)

**Table 1: Yearly count of all cases versus case count and prevalence of DM in specialist hospital Gombe (2010-2014)**

Year	2010	2011	2012	2013	2014	Total
Number of all cases (nA)	6212	5704	9710	12944	5513	40083
Number of DM cases (nD)	76	226	1219	431	117	2069
Percentage (%)	1.22	3.96	12.55	3.33	2.12	5.16

DM: Diabetes mellitus

**Table 2: DM cases by age group in specialist hospital Gombe (2010-2014)**

Age group	Years (2010-2012) N (%)	Years (2013-2014) N (%)	Total N (%)
25-34	20 (1.32)	13 (2.37)	33 (1.60)
35-44	106 (6.97)	21 (3.83)	127 (6.14)
45-54	140 (9.20)	22 (4.02)	162 (7.83)
55-64	265 (17.42)	194 (35.40)	459 (22.18)
65-75	990 (65.09)	298 (54.38)	1288 (62.25)
Total	1521 (100.00)	548 (100.00)	2069 (100.00)

$\chi^2=24.5502, df=4, P=0.000002$ . DM: Diabetes mellitus

**Table 3: Yearly distribution of diabetes mellitus cases by sex in specialist hospital Gombe (2010-2014)**

Year	2010 N (%)	2011 N (%)	2012 N (%)	2013 N (%)	2014 N (%)	Total N (%)
Male	50 (65.79)	100 (44.25)	119 (9.76)	255 (59.16)	54 (46.15)	578 (27.94)
Female	26 (34.21)	126 (55.75)	1100 (90.24)	176 (40.84)	63 (53.85)	1491 (72.06)
Total	76 (100.00)	226 (100.00)	1219 (100.00)	431 (100.00)	117 (100.00)	2069 (100.00)

$\chi^2=449.15521, df=4, P=0.0000000$

**Table 4: DM occurrence by residence in specialist hospital Gombe (2010-2014)**

Place of residence	2010 N (%)	2011 N (%)	2012 N (%)	2013 N (%)	2014 N (%)	Total N (%)
Rural	22 (28.95)	32 (14.16)	184 (15.09)	119 (27.61)	49 (41.88)	406 (19.62)
Urban	54 (71.05)	194 (85.84)	1035 (84.91)	312 (72.39)	68 (58.12)	1663 (80.38)
Total	76 (100.00)	226 (100.00)	1219 (100.00)	431 (100.00)	117 (100.00)	2069 (100.00)

$\chi^2=39.02524$ ,  $df=4$ ,  $P=0.00000$ . DM: Diabetes mellitus

while the remaining 19.62% (406/2069) resided in rural areas. This variation was quite significant with  $\chi^2 = 39.02524$ ,  $P = 0.00000$ .

## Discussion

### Overall and yearly prevalence of DM

With a computed overall DM prevalence of 5.16% and a census population figure of Gombe state (3,028,459) cited in the study setting, we can say that there is an estimated population of not <156,269 people in Gombe State that are suffering from the disease (Table 1). This rate appears to be on the high side compared to the Nigerian national prevalence rate of 4.99% according to the International Diabetes Federation (IDF).<sup>[1]</sup> When determined according to individual years covered in this study, there was a steep rise from an initial lowest rate of 1.22% in the year 2010 to 3.96% in 2011. This rising trend continued to a peak of 12.55% in the year 2012. This sharp rise in cases may have been occasioned by heightened sensitization activities by DM clubs and societies in the state at about the same period, thereby leaving the possibility of existence of an iceberg of non-reported cases of the disease in the hospital catchment communities. Thereafter, the disease prevalence nosedived to 3.33% in the year 2013 and finally to 2.12% in the year 2014. This picture contrasts with findings at global, regional and national levels where both developed and developing countries are experiencing an almost linear rising trend in the prevalence of diabetes, cardiovascular diseases, and other NCDs.<sup>[1,15]</sup> However, on the prevalence of diabetes in Nigeria, Oputa and Chinyere states in a review that, "The diabetes prevalence of 4.9% has more than doubled when compared with the 2.2% prevalence of the Professor Akinkugbe-led National Survey Report of 1997."<sup>[16]</sup> Hence, the fluctuating trend observed in this study need to be further substantiated by other more rigorous studies in the study area.

### Age pattern of DM occurrence

When stratified by age group, this study established a consistently rising pattern of DM burden with age in

which the oldest age group (65-75) was responsible for 62.25% of cases, while the lowest age rung (25-34) accounted for the smallest percentage (1.60%) of DM cases. The association between age and DM occurrence was found to be quite significant ( $\chi^2 = 24.5502$ ,  $P = 0.000002$ ). This concurs with findings of other studies from different parts of the world and within Nigeria.<sup>[10]</sup> In a study on Type 2 diabetes and its correlates among adults in Bangladesh, respondents within the age group of 55-59 years had higher odds of having diabetes (odds ratios = 2.37, 95% CI: 1.76-3.21) than the age group of 35-39 years.<sup>[17]</sup> In a study in Uyo Southern Nigeria, subjects in the highest age group in the study (46-60) were about twice more likely to have diabetes than all the others in the lower age groups.<sup>[6]</sup> In another DM prevalence study in Port Harcourt, Southern Nigeria aging was also found to be associated with Type 2 diabetes.<sup>[9]</sup>

### Sex distribution of DM cases

When segregated by sex this study showed that more females were affected than males in a ratio of 2.58:1 (72.06%/27.94%) (Table 3). This sex differential was found to be quite significant ( $\chi^2 = 449.15521$ ,  $P = 0.0000$ ). This agrees with several other studies that revealed higher female ratio in different proportions. Findings from Port Harcourt by Tamunopriye and Iroro showed 70% females and 30% males with a male-female ratio of 1:2.3.<sup>[18]</sup> Uloko *et al.* reported from a multi-center study in Nigeria where out of a total of 531 DM patients, 209 (39.4%) were males and 322 (60.6%) were females.<sup>[19]</sup> From Enugu, Southern Nigeria Chukwu *et al.* reported a higher proportion or cases of DM among females (56.0%) than the males (44.0%).<sup>[20]</sup> However, there were reports of higher male prevalence even Nigeria. In a hospital-based retrospective survey in Katsina Northern Nigeria Suleiman and Abdulkarim recorded more males that suffered from the diseases (201, 71.53%) than females (80, 28.47%).<sup>[7]</sup> A report from Abeokuta, southern Nigeria states that "Male subjects had higher prevalence of diabetes (1.85%) than females with 1.19% prevalence."<sup>[21]</sup> Omorogiuwa *et al.* also

reported higher male prevalence ratio, “The male/female diabetic ratio were 1.4:1 and 1.3:1 for senior staff and junior staff, respectively,” among university staff in Ekpoma southern Nigeria.<sup>[3]</sup> A sedentary lifestyle of women in this part of the country could be a possible explanation to our findings of a feminine preponderance of DM case burden. Men are adjudged to be more physically active than women, and this probably could enhance insulin sensitivity in the men than women. Ekpenyong *et al.* and Nyenwe *et al.* found physical inactivity to be significantly associated with the incidence of diabetes in their study.<sup>[6,9]</sup> The male preponderance in other studies may not reflect the real situation because the pattern of hospital medical admissions may also need to be examined in such studies. The male bias may, therefore, be related to cultural, geographic and socioeconomic factors rather than the gender-related prevalence of DM.<sup>[21]</sup>

#### Monthly trend of DM cases

Cumulative monthly trend of cases within the study period (2010-2014) exhibited an overall rising pattern of DM case occurrence from January to December. Three remarkable peaks were recorded - in July (141 cases), October (428 cases) and December (560 cases), while two troughs were observed in August (61 cases) and November (337 cases). October and December peaks were more pronounced, suggesting a seasonal predilection of DM case occurrence toward the end of raining season and the peak of harvest and festive period of the year. This tendency for having high reports of cases of the disease within these periods could be due to the high likelihood of locals breaching dietary control habits during festivities and harvest seasons when varieties of food abound. A seasonality of Type 1 diabetes has been reported in Tanzania, with most cases presenting between August and November.<sup>[22]</sup> From the University of Benin Teaching Hospital, Benin City, Nigeria, Edo *et al.* reported peak cases in July and December, with the highest in December.<sup>[23]</sup> In a 6 years study on, “Seasonal Incidence of DM in Tropical Africa” the peak months of presentation were August through to November, with the greatest number of patients presenting in September.<sup>[24]</sup> In a slightly varied scenario Tamunopriye and Iroro from Port Harcourt reported in their study that, “67% of patients were diagnosed in the rainy season.”<sup>[18]</sup>

#### Prevalence of DM by place of residence

This study recorded more urban (1663, 80.38%) than rural (406, 19.62%) cases of DM. The differences

were statistically significant ( $\chi^2 = 39.02524$ ,  $P = 0.00000$ ). This agrees with the majority of findings at various levels. A retrospective hospital-based study in Katsina, Northern Nigeria, established an overwhelmingly higher case burden in the urban (82.56%) than rural (17.44%) areas.<sup>[7]</sup> In Lagos South Western Nigeria, Oguoma *et al.* also reported higher urban (5.4%) than rural (1.8%) prevalence figures of DM.<sup>[25]</sup> Moreover, the explanation to this discrepancy has consistently been fast changing lifestyle due to the overwhelming influence of westernization cum urbanization in most developing countries. “Urbanization tends to decrease physical activity as very little physical activity is required for daily living whereas rural populations rely on foot walk as transportation means and often have intense agricultural activities as their main occupation.”<sup>[11]</sup>

#### Conclusion

The findings of this study show clearly that the prevalence of DM in the study area is consistent with the national prevalence according to the IDF. However, the yearly trend which exhibited a sharp rise in the year 2012 and a nosedive thereafter might be indicative of some dramatic progress in the control efforts occasioned by massive media and community sensitization programs on appropriate lifestyle modifications. In any case, the disproportionate distribution of the disease burden with obvious bias toward females, urban dwellers and the aged are useful information for the design of more targeted prevention programs by the government and other stakeholders.

#### References

1. International Diabetes Federation. Global Diabetes Scorecard Tracking Progress for Action. Oman: IDF; 2014. p. 5-24.
2. Akinjinmi AA, Adeyooye OA, Akingbade OA, Okerentugba PO. Prevalence of Diabetes mellitus in Abeokuta, Ogun state, Nigeria. Researcher 2014;6:73-5. Available from: <http://www.sciencepub.net/researcher>.10. [Last accessed on 2015 Dec 05].
3. Omorogiuwa A, Oaikhena GA, Okioya P, Akubueze D, Owobu E, Enahoro I, *et al.* Diabetes mellitus: Prevalence amongst university staff in Southern Nigeria and attitude towards routine glycaemic/glucosuric checkup. Int J Biomed Health Sci 2010;6:25-9.
4. Ejike CE, Uka NK, Nwachukwu SO. Diabetes and pre-diabetes in adult Nigerians: Prevalence, and correlations of blood glucose concentrations with measures of obesity. Afr J Biochem Res 2015;9:55-60.
5. Isara AR, Okundia PO. The burden of hypertension and diabetes mellitus in rural communities in southern Nigeria. Pan Afr Med J 2015;20:103.
6. Ekpenyong CE, Akpan UP, Ibu JO, Nyebuk DE. Gender and age specific prevalence and associated risk factors of Type 2

- diabetes mellitus in Uyo Metropolis, South Eastern Nigeria. *Diabetol Croat* 2012;41:17-28.
7. Suleiman M, Abdulkarim B. Incidence of diabetes mellitus at the Katsina General Hospital, Katsina State, Nigeria. *Appl Sci Rep* 2015;11:29-32. DOI: 10.15192/PSCP.ASR.2015.11.1.2932. Available from: <http://www.pscipub.com>. [Last accessed on 2015 Dec 05].
  8. National Population Commission, (NPC). Nigeria National Census, 2006. Available from: [http://www.population.gov.ng/pop\\_figure.pdf](http://www.population.gov.ng/pop_figure.pdf). [Last accessed on 2015 Dec 05].
  9. Nyenwe EA, Odia OJ, Ihekwa AE, Ojule A, Babatunde S. Type 2 diabetes in adult Nigerians: A study of its prevalence and risk factors in Port Harcourt, Nigeria. *Diabetes Res Clin Pract* 2003;62:177-85.
  10. Olokoba AB, Obateru OA, Olokoba LB. Type 2 diabetes mellitus: A review of current trends. *Oman Med J* 2012;27:269-73.
  11. Hall V, Thomsen RW, Henriksen O, Lohse N. Diabetes in Sub Saharan Africa 1999-2011: Epidemiology and public health implications. A systematic review. *BMC Public Health* 2011;11:564.
  12. Dahiru T, Jibo A, Hassan AA, Mande AT. Prevalence of diabetes in a semi-urban community in Northern Nigeria. *Niger J Med* 2008;17:414-6.
  13. Sabir A, Ohwovoriole A, Isezuo S, Fasanmade O, Abubakar S, Iwuala S. Type 2 diabetes mellitus and its risk factors among the rural Fulanis of Northern Nigeria. *Ann Afr Med* 2013;12:217-22.
  14. Bakari AG, Onyemelukwe GC, Sani BG, Hassan SS, Aliyu TM. Prevalence of diabetes in suburban northern Nigeria: Results of a public screening survey. *Diabetes Int* 1999;9:59-60.
  15. Jayawardena R, Ranasinghe P, Byrne NM, Soares MJ, Katulanda P, Hills AP. Prevalence and trends of the diabetes epidemic in South Asia: A systematic review and meta-analysis. *BMC Public Health* 2012;12:380.
  16. Oputa RN, Chinenye S. Diabetes mellitus: A global epidemic with potential solutions. *Afr J Diabetes Med* 2012;20:33-5.
  17. Chowdhury MA, Uddin MJ, Khan HM, Haque MR. Type 2 diabetes and its correlates among adults in Bangladesh: A population based study. *BMC Public Health* 2015;15:1070.
  18. Tamunopriye J, Iroro Y. The pattern of presentation and trends of childhood diabetes mellitus in Port Harcourt, Southern Nigeria. *Br J Med Med Res* 2015;5:247-53.
  19. Uloko AE, Ofoegbu EN, Chinenye S, Fasanmade OA, Fasanmade AA, Ogbera AO, *et al.* Profile of Nigerians with diabetes mellitus - Diabcare Nigeria study group (2008): Results of a multicenter study. *Indian J Endocrinol Metab* 2012;16:558-64.
  20. Chukwu BN, Ezebuio VO, Samuel ES, Nwachukwu KC. Gender differential in the incidence of diabetes mellitus among the patients in UDI local government Area of Enugu State, Nigeria. *Mediterr J Soc Sci* 2013;4:131-8.
  21. Abdulkadir J. Malnutrition-related diabetes mellitus in Africa. *Int J Diabetes Dev Ctries* 1993;13:22-8.
  22. Mbanya JC, Ramiya K. Diabetes mellitus. Disease and Mortality in Sub-Saharan Africa. 2<sup>nd</sup> ed. Ch. 19. Brussels: IDF; 2006.
  23. Edo AE, Edo GO, Ezeani IU. Risk factors, ulcer grade and management outcome of diabetic foot ulcers in a Tropical Tertiary Care Hospital. *Niger Med J* 2013;54:59-63.
  24. McLarty DG, Yusafali A, Swai AB. Seasonal incidence of diabetes mellitus in tropical Africa. *Diabet Med* 1989;6:762-5.
  25. Oguoma VM, Nwose EU, Skinner TC, Digban KA, Onyia IC, Richards RS. Prevalence of cardiovascular disease risk factors among a Nigerian adult population: Relationship with income level and accessibility to CVD risks screening. *BMC Public Health* 2015;15:397.

**Financial Support:** None; **Conflict of Interest:** None

**How to cite this article:** Danjin M, Umar NU, Adamu D. Prevalence of diabetes mellitus in a tertiary health institution in Gombe Metropolis, Nigeria. *J Med Sci Health* 2017;3(1):1-6.

Date of submission: 09-06-2016

Date of review: 29-11-2016

Date of acceptance: 04-12-2017

Date of publication: 10-03-2017