

Impact of Linear Vestibular Stimulation on Lipid Profile in Underweight Females: A Pilot Study

S Saritha¹, Kumar Sai Sailesh², J K Mukkadan³

ABSTRACT

Background and Objectives: Weight gain and weight loss form complex health issues. Animal studies provided preliminary evidence for the regulation of total cholesterol and TG through vestibular stimulation. However, human studies on the same area are sparse. Hence, the present study was aimed to assess the impact of vestibular stimulation on lipid profile in underweight females.

Materials and Methods: A total of 8 underweight females and 8 healthy, age-matched, females were included in the present study after obtaining voluntary informed consent. Vestibular stimulation was administered by making the participants to swing on a swing in back-to-front direction. Total cholesterol, triglycerides, high-density lipoprotein (HDL), and low-density lipoprotein (LDL) were estimated by standard methods.

Results: Total cholesterol and LDL levels were significantly lower in underweight females before and after the intervention ($P < 0.001$). HDL and triglycerides levels are significantly lower ($P < 0.001$) in underweight females before the intervention. The difference was not statistically significant after the intervention.

Conclusion: We have observed an increase in triglycerides and HDL and no change in LDL and total cholesterol followed by vestibular stimulation. We recommend further detailed studies in this area for a better understanding of vestibular influences on lipid profile which may help to develop a simple, novel supplementary therapy in the management of metabolic disorders.

KEY WORDS: Linear vestibular stimulation, lipid profile, underweight females.

Introduction

According to the WHO definition, BMI is the simple index of weight for height that commonly used to classify overweight and obesity in adults.^[1] BMI is defined as the person weight in kilogram divided by the square of the height in m².^[2] Intake of nutrients is under complex control involving signal from central nervous system and peripheral nervous system. Vestibular system having extensive interaction between the hypothalamus, dorsal raphe nucleus, and tractus solitarius regulates food intake.^[3] Weight gain and weight loss form complex health issues. The current animal study observed that vestibular stimulation can effectively regulate total cholesterol and TG which helps to maintain healthy body

weight.^[4] Animal studies provided preliminary evidence for the regulation of total cholesterol and TG through vestibular stimulation. However, human studies on the same area are sparse. Hence, the present study was aimed to assess the impact of vestibular stimulation on lipid profile in underweight females.

Materials and Methods

The present experimental study was conducted at the Department of Physiology, Little Flower Institute of Medical Sciences and Research, Angamaly.

Participants

A total of 8 underweight females and 8 healthy, age-matched, females were included in the present study after obtaining voluntary informed consent. The following criteria were used for recruiting the underweight females.

Inclusion criteria

The following inclusion criteria were included in the study:

1. Apparently healthy and willing participants,

Access this article online

Quick Response Code:



Website: www.jmsh.ac.in

¹Post Graduate Student, Department of Physiology, Little Flower Institute of Medical Sciences and Research, Angamaly, Kerala, India, ²Assistant Professor, Department of Physiology, D M Wayanad Institute of Medical Sciences, Wayanad, Kerala, India, ³Research Director, Little Flower Institute of Medical Sciences and Research, Angamaly, Kerala, India

Address for correspondence:

Sai Sailesh kumar G, Assistant Professor, Department of Physiology, DM Wayanad Institute of Medical Sciences, Naseera Nagar, Meppadi, Wayanad, Kerala, India. E-mail: saisailesh.kumar@gmail.com

2. Underweight female students (BMI <18.50)
3. Age of 18-24 years.

Exclusion criteria

The following exclusion criteria were excluded from the study:

1. Participants with hearing, vestibular disorders,
2. Those under any medications including oral contraceptives
3. Participants under any kind of treatment or therapy.
 - Group A: ($n = 8$) healthy normal weight students without intervention
 - Group B: ($n = 8$) underweight students without intervention
 - Group C: ($n = 8$) underweight students with intervention for 60 days.

After recording baseline values in all the groups, vestibular stimulation was administered for intervention groups for 60 days. Post-intervention values were recorded in all the groups and compared.

Methods

Linear vestibular stimulation

Vestibular stimulation was administered by making the participants to swing on a swing in back-to-front direction (according to comfort), as standardized by the previous methods.^[5-7]

Outcome measures

Total cholesterol was estimated by cholesterol oxidase-phenol+aminophenazone (CHOD PAP) method, triglycerides was estimated by glycerol-3-phosphate oxidase-phenol+aminophenazone (GPO-PAP) method, and high-density lipoprotein (HDL) and low-density lipoprotein (LDL) were estimated by precipitation method.

Data analysis

Data were analyzed by SPSS 20.0. Unpaired t-test was used to observe the significance of difference in demographic data. Two-way ANOVA followed by Bonferroni post-test was used to observe the significance of difference in our come measures.

Results

Figure 1 presents demographic data of the participants. Body weight and BMI was significantly lower in underweight females when compared with healthy females. Height was not significantly different. Figure 2 presents HDL, LDL, triglycerides,




Figure 1: Demographic data of the participants (data were expressed as mean \pm SD). * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$

and total cholesterol of the participants. Total cholesterol and LDL levels were significantly lower in underweight females before and after the intervention ($P < 0.001$). HDL and triglycerides levels are significantly lower ($P < 0.001$) in underweight females before the intervention. The difference was not statistically significant after the intervention.

Discussion

The current study was undertaken to evaluate the impact of vestibular stimulation on lipid profile. We have observed an increase in the triglycerides and HDL followed by vestibular stimulation. When compared with healthy controls, the baseline values of these parameters were significantly lower before the intervention. Followed by vestibular stimulation, there was an increase in the triglycerides and HDL, so the difference was not statistically significant when compared with control group after the intervention. No significant change was observed in total cholesterol and LDL. Earlier studies reported that vestibular stimulation regulates taste aversion and food intake.^[8-14] Vestibular stimulation was reported to modulate the biochemical parameters.^[15-18] Earlier animals study applied caloric vestibular stimulation in an animal model of hyperlipidemia. It was reported that there was a significant decrease in total cholesterol and triglycerides and a significant decrease in the HDL and LDL.^[4] In the present study, vestibular stimulation was administered using a swing, and we have observed an increase in the triglycerides and HDL. It may be due to the difference in the type of stimulation that influences the lipid profile.




Figure 2: High-density lipoprotein, low-density lipoprotein, triglycerides, and total cholesterol of the participants. (data were expressed as mean ± SD) **P* < 0.05, ***P* < 0.01, ****P* < 0.001

Limitations

The major limitation of our study was less sample size.

Conclusion

We have observed an increase in triglycerides and HDL and no change in LDL and total cholesterol followed by vestibular stimulation. We recommend further detailed studies in this area for a better understanding of vestibular influences on lipid profile which may help to develop a simple, novel supplementary therapy in the management of metabolic disorders.

References

1. WHO. Physical status: The use and interpretation of anthropometry. Report of a WHO Expert Committee. WHO Technical Report Series 854. Geneva: World Health Organization; 1995.
2. WHO Expert Consultation. Appropriate body-mass index for Asian populations and its implications for policy and intervention strategies. *Lancet* 2004;363:157-63.
3. Sailesh KS. Vestibular balance of food intake. *Int J Pharm Bio Sci* 2014;5:1069-73.
4. Sadanandan NN, Archana R, Sailesh KS, Antony NJ. Antihyperlipidemic effect of vestibular stimulation in wistar albino rats. *Int J Res Ayurveda Pharm* 2015;6:509-12.
5. Sailesh KS, Archana R, Antony NJ, Mukkadan JK. You

- are never too old to swing. *Res J Pharm Biol Chem Sci* 2014;5:612-5.
6. Sailesh KS, Mukkadan JK. Controlled vestibular stimulation, standardization of a physiological method to release stress in college students. *Indian J Physiol Pharmacol* 2015;59:436-41.
7. Sailesh KS Archana R, Mukkadan JK. Controlled vestibular stimulation: A physiological method of stress relief. *J Clin Diagn Res* 2014;8:BM01-2.
8. Brozek G. Electrophysiological analysis of conditioned taste aversion in rats. *Acta Neurobiol Exp (Wars)* 1982;42:29-41.
9. Laskiewicz J, Królczyk G, Zurowski G, Sobocki J, Matyja A, Thor PJ. Effects of vagal neuromodulation and vagotomy on control of food intake and body weight in rats. *J Physiol Pharmacol* 2003;54:603-10.
10. Gil K, Bugajski A, Thor P. Electrical vagus nerve stimulation decreases food consumption and weight gain in rats fed a high-fat diet. *J Physiol Pharmacol* 2011;62:637-46.
11. Filippi BM, Bassiri A, Abraham MA, Duca FA, Yue JT, Lam TK. Insulin signals through the dorsal vagal complex to regulate energy balance. *Diabetes* 2014;63:892-9.
12. Silverstein JT, Plisetskaya EM. The effects of NPY and insulin on food intake regulation in fish. *Am Zool* 2000;40:296-308.
13. Seicean A, Redline S, Seicean S, Kirchner HL, Gao Y, Sekine M, *et al.* Association between short sleeping hours and overweight in adolescents: Results from a US Suburban High School survey. *Sleep Breath* 2007;11:285-93.
14. Alva-Sánchez C, Pacheco-Rosado J, Fregoso-Aguilar T, Villanueva I. The long-term regulation of food intake and body weight depends on the availability of thyroid hormones in the brain. *Neuro Endocrinol Lett* 2012;33:703-8.

15. Kekhaiov AN. Sugar, chlorides and cholesterol in the blood of pilots before and after vestibular stimulation. Zh Ushn Nos Gorl Bolezn 1976;5:21-4.
16. Kekhaiov AN. Various serum biochemical indices in patients with vestibular disorders. Zh Ushn Nos Gorl Bolezn 1971;31:52-5.
17. Kekhaiov AN. Biochemical changes in the serum of patients with pronounced vestibulo-autonomic and vestibulo-somatic reactions. Zh Ushn Nos Gorl Bolezn 1971;31:67-8.
18. Fuller P, Warden C, Barry S, Fuller C. Effects of 2-G exposure on temperature regulation, circadian rhythms, and adiposity in UCP2/3 transgenic mice. J Appl Physiol 2000;89: 1491-8.

Financial Support: None; **Conflict of Interest:** None

How to cite this article: Saritha S, Sailesh KS, Mukkadan JK. Impact of Linear Vestibular Stimulation on Lipid Profile in Underweight Females: A Pilot Study. J Med Sci Health 2017;3(3):43-46.

Date of submission: 05-04-2017

Date of review: 26-08-2017

Date of acceptance: 17-10-2017

Date of publication: 15-12-2017